

VU Research Portal

MERIS-based ocean colour classification with the discrete Forel-Ule scale

Wernand, M.R.; Hommersom, A.; van der Woerd, H.J.

published in
Ocean Science
2013

DOI (link to publisher)
[10.5194/os-9-477-2013](https://doi.org/10.5194/os-9-477-2013)

document version
Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Wernand, M. R., Hommersom, A., & van der Woerd, H. J. (2013). MERIS-based ocean colour classification with the discrete Forel-Ule scale. *Ocean Science*, 9, 477-487. <https://doi.org/10.5194/os-9-477-2013>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:
vuresearchportal.ub@vu.nl

This discussion paper is/has been under review for the journal Ocean Science (OS).
Please refer to the corresponding final paper in OS if available.

MERIS-based ocean colour classification with the discrete Forel–Ule scale

M. R. Wernand¹, A. Hommersom², and H. J. van der Woerd²

¹Royal Netherlands Institute for Sea Research, Physical Oceanography, Marine Optics & Remote Sensing, P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands

²Institute for Environmental Studies (IVM), VU University Amsterdam, De Boelelaan 1087, 1081 HV Amsterdam, The Netherlands

Received: 13 July 2012 – Accepted: 16 August 2012 – Published: 31 August 2012

Correspondence to: M. R. Wernand (marcel.wernand@nioz.nl)

Published by Copernicus Publications on behalf of the European Geosciences Union.

OSD

9, 2817–2849, 2012

MERIS-based Forel–Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Multispectral information from satellite borne ocean colour sensors is at present used to characterize natural waters via the retrieval of concentration of the three dominant optical constituents; pigments of phytoplankton, non-algal particles and coloured dissolved organic matter. A limitation of this approach is that accurate retrieval of these constituents requires detailed local knowledge of the specific absorption and scattering properties. In addition, the retrieval algorithms generally use only a limited part of the collected spectral information. In this paper we present an additional new algorithm that has the merit to use the full spectral information in the visible domain to characterize natural waters in a simple and globally valid way. This Forel–Ule MERIS (FUME) algorithm converts the normalized multi-band reflectance information into a discrete set of numbers using uniform colourimetric functions. The Forel–Ule scale is a sea colour comparator scale that has been developed to cover all possible natural sea colours, ranging from indigo blue (the open ocean) to brownish-green (coastal water) and even brown (humic-acid dominated) waters. Data using this scale have been collected since the late nineteenth century, and therefore, this algorithm creates the possibility to compare historic ocean colour data with present-day satellite ocean colour observations. The FUME algorithm was tested by transforming a number of MERIS satellite images into Forel–Ule colour index images and comparing in situ observed FU numbers with FU numbers modelled from in situ radiometer measurements.

1 Introduction

The application of optical satellite remote sensing techniques to monitor the radiation scattered back from the water column became a major breakthrough in the seventies for monitoring ocean, sea and coastal areas (IOCCG, 1998). Dedicated ocean colour instruments, like CZCS, SeaWiFS, MERIS and MODIS-AQUA, have provided fundamental new insight in the dynamics and role of oceanic plankton (e.g. Behrenfeld et al.,

OSD

9, 2817–2849, 2012

MERIS-based Forel–Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2006). Observations are now starting to span multiple decades, allowing a first glimpse at long-term variations in the plankton composition of the oceans, which are potentially related to global change (Antoine et al., 2005; Polovina et al., 2008).

With the launch, in 2002, of the MERIS instrument (Rast et al., 1999) that measures water-leaving reflectance in fifteen spectral bands with high signal-to-noise, it became possible to collect water-leaving radiance with high confidence in regional seas and coastal waters. This has led to development of many new algorithms that can retrieve not only the phytoplankton pigments, but also the mass concentration of suspended material and the absorption by dissolved material (Van der Woerd and Pasterkamp, 2008; Odermat et al., 2012). These algorithms are either simple, calibrated at the local water constituents, or complex with a need for detailed measurements of the specific absorption and scattering properties of these in-water constituents (see e.g. Tilstone et al., 2012). The derived water-quality parameters are the major products of ocean-colour instruments, while the colour itself can be considered as a primary product.

Long before the development of diode-arrays to measure spectral radiation, another method had been developed and tested which recorded the colour of natural waters. Towards the end of the nineteenth century, Forel and Ule (Forel, 1890; Ule, 1892) proposed a method to classify the colour of the oceans, regional seas and coastal waters using a colour comparator scale. The scale became known as the Forel–Ule (FU) scale and since then scale observations have been performed, generating hundreds of thousands of data points over a globe-scale for more than a century. Recently, it was shown (Wernand and Van der Woerd, 2010a) that the FU scale is can be used to characterize the colour of natural waters. More importantly, the analysis of FU colour variation in the North Pacific since 1930 has revealed significant variations at decadal timescales (Wernand and Van der Woerd, 2010b).

In this paper we describe a simple algorithm to couple historically collected ocean colour data, obtained over a long time span, with presently available satellite-derived ocean colour imagery for hind casting long-term changes. This Forel–Ule to MERIS (FUME) algorithm converts MERIS observations of sea- and ocean colour to

MERIS-based Forel-Ule classification

M. R. Wernand et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

chromaticity coordinates and subsequently to a discrete Forel–Ule number. This will result in a new MERIS water quality product that can be used as a simple and straightforward index of water colour in addition to the water-quality parameters that are retrieved by inversion schemes. Based on the FUME product, ocean colour trends can be constructed, reaching back to over one hundred years.

2 Methods

In this section we introduce the MERIS satellite data, the algorithm to convert MERIS reflection data to FU numbers and the ship-borne measurements for a first characterisation of the FUME results.

2.1 MERIS products

MERIS is a 68.5° field-of-view push-broom imaging spectrometer (Rast et al., 1999) on the ENVISAT platform. It measures the solar radiation reflected by the ocean at a spatial resolution of 260 m × 290 m in 15 spectral bands. The bands are programmable in width and position, at visible and near infrared wavelengths. MERIS provides global coverage in 3 days with radiation reflected by the ocean that is atmospherically corrected to derive the normalized water-leaving reflectances, a MERIS Level 2 product (ESA, 2012). The atmospheric correction assumes that the water totally absorbs the NIR, but also includes a correction for those sediment loaded waters where this assumption fails. Normalized water-leaving reflectance (dimensionless) $[\rho_W]_N$ is defined by Eq. (1) in as:

$$[\rho_W]_N(\lambda) = \frac{[L_W]_N(\lambda)}{F_0(\lambda)} \quad (1)$$

where $[L_W]_N$ is the normalized water-leaving radiance (Gordon and Voss, 1999) and F_0 is the extraterrestrial solar irradiance at wavelength (λ). In this analysis, data is limited

MERIS-based Forel–Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the visible spectrum, covering the first nine MERIS bands with a bandwidth of 10 nm, except for Band 8 that has a bandwidth of 7.5 nm (Table 1). In the standard processing by ESA a number of global products are derived together with $[\rho_W]_N$ that will be used to compare FU products with standard ESA products: Algal-1 and Algal-2, the index for Chlorophyll *a* concentration in case-1 and case-2 waters, respectively, SPM (suspended particulate matter) and YS (an index for absorption by dissolved matter). For documentation and additional references we recommend the MERIS algorithm theoretical baseline document (ESA, 2012).

2.2 The FUME algorithm

The FUME algorithm converts the normalized water-leaving reflectance in nine MERIS bands into a discrete FU number in three steps: Step 1: Calculation of the tristimulus values X , Y , Z by calculating the convolution of the Colour Matching Functions (CMF's) and the normalized water-leaving reflectance (CIE, 1932). Step 2: Calculation of the (x, y) chromaticity coordinates by the ratio of X or Y tristimulus value and the sum of the tristimulus values. Step 3: Determination of the FU scale number by comparison of calculated (x, y) values to the unique chromaticity coordinates of the twenty-one FU numbers.

Step 1: Tristimulus values are the amounts of three primaries that specify a colour stimulus of the human eye (Wyszecky and Stiles, 2000) and are noted as X , Y and Z (CIE, 1932). The CIE 1931 standard colourimetric 2 degree CMFs \bar{x} (red), \bar{y} (green) and \bar{z} (blue) are presented in Fig. 1. These serve as weighting functions for the determination of the tristimulus values of the MERIS normalized water-leaving reflectance

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$[\rho_W]_N$ by the Eqs. (2a), (2b) and (2c);

$$X = \int [\rho_W]_N(\lambda) \bar{x}(\lambda) d\lambda \quad (2a)$$

$$Y = \int [\rho_W]_N(\lambda) \bar{y}(\lambda) d\lambda \quad (2b)$$

$$Z = \int [\rho_W]_N(\lambda) \bar{z}(\lambda) d\lambda \quad (2c)$$

Because MERIS does not provide full-spectral coverage, the reflection spectrum is first reconstructed by linear interpolation between band $n = 1$ (412.5 nm) and band $n = 9$ (708 nm) with a resolution of 1 nm. An example is shown as black line in Fig. 1. Note that the linear interpolation at λ_i (nm) is always carried out between subsequent bands ($n, n + 1$) with the condition ($\lambda_n < \lambda_i < \lambda_{n+1}$). The tristimulus values for X , Y and Z are obtained by a Riemann sum approximation of the integrals with $\Delta\lambda = 1$ nm resolution;

$$X = \sum_{i=413}^{708} [\rho_W]_N(\lambda_i) \bar{x}(\lambda) \Delta\lambda \quad (3a)$$

$$Y = \sum_{i=413}^{708} [\rho_W]_N(\lambda_i) \bar{y}(\lambda) \Delta\lambda \quad (3b)$$

$$Z = \sum_{i=413}^{708} [\rho_W]_N(\lambda_i) \bar{z}(\lambda) \Delta\lambda \quad (3c)$$

Step 2: Subsequently the chromaticity coordinates x , y and z are calculated from the ratio of each of the tristimulus values and the sum of the values:

$$x = \frac{X}{X+Y+Z} \quad y = \frac{Y}{X+Y+Z} \quad z = \frac{Z}{X+Y+Z} \quad (4)$$

As $x + y + z = 1$, and therefore $z = 1 - x - y$, the third coordinate offers no additional information and only two coordinates (by convention x and y) are used to represent the colour in a so-called chromaticity diagram (see e.g. Mobley, 1994). The white point

W in the chromaticity diagram has the coordinates $x = y = z = 1/3$ (Fig. 2). The ratio of the distance between W and an arbitrary point P (a) and the distance from W to the spectral locus ($a + b$), gives the colour saturation ($a/(a + b)$) or the intensity of the colour in P. In this way, the chromaticity coordinates (x_M, y_M) for every MERIS pixel can be calculated.

Step 3: In the next step the (x_M, y_M) is converted to a FU number. The original FU scale was created to make an objective classification of natural waters (see for a review Wernand and Gieskes, 2011). In 21 glass tubes a variable mixture of three standard solutions (distilled water, ammonia, copper-sulphate, potassium-chromate and cobalt-sulphate) were created to obtain the colour-palettes of the scale. These standard solutions were recently reconstructed and their optical properties were measured in the laboratory with medium resolution spectrometers (Wernand and Van der Woerd, 2010a). The calculated chromaticity coordinates of the original FU scale are presented in Table 2 and graphically shown as a line of black dots, between the white point and the locus, in the chromaticity diagram of Fig. 2. The FUME algorithm first shifts the origin to the white point W with chromaticity coordinates $x_W = y_W = 1/3$ (Fig. 3). Then it calculates the angle (α_M) between the vector to a point with certain FU coordinates (x_M, y_M) and the positive x-axis (at $y - y_W = 0$), giving higher angles in an anti-clockwise direction and compares these with the angles (α_i) of the FU solutions (Table 3). All calculations were made with the ATAN2 function (four-quadrant inverse tangent) and the derived angles (in radials) were multiplied by $180/\pi$ to get the angles in degrees:

$$\alpha_M = \arctan(y_M - y_W, x_M - x_W) \text{ modulus } 2\pi \quad (5)$$

Two examples are shown in Fig. 3: α_i is the angle matching FU number 8. The yellow dot is derived from the normalized spectral reflectance of a MERIS-pixel and coordinates ($x_M - x_W = -0.15$, $y_M - y_W = 0.1$) and angle (α_M). Finally, the boundaries distinguishing the various FU numbers were defined. The colour transition angle α_{iT} , under

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which a scale number transition takes place, was taken according to Eq. (6).

$$\alpha_{iT} = \frac{(\alpha_i + \alpha_{i+1})}{2} \quad (6)$$

Both α_i and α_{iT} are presented in Table 3. The FU numbers for a given MERIS pixel M with chromaticity coordinates $x_M - x_W = -0.15$ and $y_M - y_W = 0.1$ (yellow point in Fig. 3) can be determined as follows: first the angle (Eq. 5) is determined as $\alpha_M = 146^\circ$ and then is compared with a simple MATLAB loop for $i = 1$ to 21 twenty-one values of α_{iT} given in Table 3. From this loop $\alpha_M > \alpha_{iT}$ is true for the first time reaching the angle $\alpha_{iT} = 133.96$ degrees that corresponds to a discrete value of FU = 6 that is attributed to this MERIS pixel M.

2.3 Ship-borne measurements

Next to the North Sea and the Wadden Sea (Hommersom et al., 2009), a shallow coastal area bordering the North Sea north of Holland and Germany (Fig. 5), rivers and lakes were optically sampled in 2001, 2006 and in 2007. The surface radiance L_{sfc} , sky radiance L_{sky} and incoming solar irradiance E_s were measured simultaneously, using TRIOS hyper-spectral radiometers (Heuermann et al., 1999). Remote sensing reflectance was then calculated as $R_{\text{RS}} = L_w/E_s$, where L_w is the water-leaving radiance ($= L_{\text{sfc}} - \rho L_{\text{sky}}$) and E_s is the downward irradiance just above the sea surface (Mueller et al., 2003). To a good approximation $[\rho_w]_N \approx \pi R_{\text{RS}}$ (Lee et al., 1994).

To illustrate the potential use of the satellite derived FU maps, databases containing globally collected ship-borne FU observations were consulted. From the oceanographic and meteorological database, archived by the United States National Oceanographic Data Centre (NOAA-NODC; Boyer et al., 2006) and from the ocean colour database at the Royal Netherlands Institute for Sea Research, FU observations were extracted. To create the maps, FU measurements were interpolated through an Inverse Distance Weighted (IDW) technique (Watson and Philip, 1985) in an ARCGIS environment. The IDW interpolation was carried out over 2 degrees with a grid size of 0.2 degrees.

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Data sets

The FUME algorithm was applied to five MERIS images acquired over the areas shown in Fig. 4. These areas were chosen for their different sea colour properties (Wernand et al., 2012) and cover the North Sea (1), the Red Sea (2, 3), Yellow Sea (4) and the Sea of Japan (5). The images were extracted from ESA's on-line database, the MERIS Catalogue and Inventory (MERCI, Brockman et al., 2005). MERIS Reduced Resolution (RR) geophysical products (Table 4) contain, among other products, a total of 14 spectral images of normalized band reflectances and the derived products for pigments (Algal-1, Algal-2, SPM and YS). A Reduced Resolution image has 4×4 less pixels than the same image in Full-Resolution, thus representing an area of $1040 \text{ m} \times 1160 \text{ m}$.

To validate the FUME algorithm a dataset of simultaneously collected FU-observations and hyperspectral subsurface and above water spectra was consulted. This dataset was established in 2001 and contains observations and optical data of a wide range of coloured water types, as river-, lake-, coastal and open sea, with the FU-scale varying from FU3 to FU21. From the 156 data sets collected in the Wadden Sea (Table 2 in Hommersom et al., 2009) only one was collected close (2.5 h prior) to the MERIS image acquisition time on 4 May 2006.

The routine collection of FU-measurements in all world seas has once been very intense, mainly in the 20th century, and over 220 000 measurements are known and available (Wernand et al., 2012). However, in the first decade of the 21st century the FU data collection is much more limited and/or has not yet been recorded in the central archives. Therefore, we have chosen to show data from the same seasons in earlier years: For the Red Sea, 52 observations are available and were collected during the winters of 1895 to 1898. For the Yellow Sea, 2882 FU observations were collected during the winters of 1930 to 1999.

OSD

9, 2817–2849, 2012

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Results

4.1 MERIS FU maps

For all five MERIS images the reflectance values in band 1–9 per pixel were converted to chromaticity coordinates and into a FU number using Eqs. (3) to (5). Converted images are further referred to as FU maps. These FU maps are presented in Figs. 5, 6 and 7. In these figures we have used the MERIS flags per pixel to identify land (grey), clouds (white) and the failure to collect observations or retrieve water-leaving reflectances (black). The legend (RGB values are given in Table 5) represents the FU colours as close as possible. The $[\rho_W]_N$ spectral signatures at the locations marked with a red circle in the maps are plotted in Fig. 8.

The first FU map shown in Fig. 5, acquisition date of 4 May 2006, covers the North Sea, the Baltic and the Wadden Sea. The colour of the North Sea varies between FU3 and FU9. The colour within the left red circle situated between the Thames and Humber estuaries was estimated as FU6. The Central North Sea shows values of FU3 to FU4 with an occasionally FU2 (very blue oceanic waters). The Wadden Sea, a large intertidal sea behind multiple barrier islands, north of Holland and Germany, and west of Denmark, is dominated by sediment and outflow of humic-acid rich river water and has higher FU values, up to FU = 18.

Figure 6 shows a winter FU map of both Northern- and Southern Red Sea taken on 22 and 23 December 2003, respectively. The colour of the Northern Red Sea is mainly FU2 to FU3 with maximum values of FU5. The southern part, which is shallower than the northern part, shows a possible plankton bloom starting south of 17.5° N with values of FU8 (red circle) to FU11. The water flowing through the narrow strait of Bab-al-Mandab into the Gulf of Aden (area at the most south-eastern point on the map) shows much bluer values: FU2 to FU4.

Figure 7 shows a FU map of the Yellow Sea (left) and the Sea of Japan (right) acquired on 11 February 2009 and 14 June 2004, respectively. The outflow of the Yangtze River (south of the red circle in the left image) shows high FU values, between

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



FU7 up to real brownish colours of FU19. Unfortunately, the area close to the river outflow is flagged as “No Data area”. Within the red circle a value of FU9 is calculated. The Sea of Japan (right panel) shows values of FU2 to FU3 (within the red circle the value is FU2). Remarkable is the relative green area east of Hokkaido (FU9 marked by the red circle east) with values up to FU10. To verify our results the MERIS level2 chlorophyll product was consulted, which showed high concentrations of chlorophyll *a* ($> 2 \text{ mg m}^{-3}$) East of Hokkaido and concentrations between 0.1 and 0.5 mg m^{-3} in the Sea of Japan.

4.2 Ground truth

The reflection spectrum at the match-up station in the Wadden Sea (WS-GT) is plotted in Fig. 8 and appears very similar in shape to the MERIS spectrum (WS). By extraction of the reflection at exactly all 9 MERIS bands and running the FUME algorithm a value of $\text{FU} = 15$ was retrieved. The MERIS pixel at this location (red circle in the Wadden Sea) has a calculated FU value of 14, which is in good agreement with the ground truth FU value concerning possible adjacency effects of tidal flats within the pixel.

The MERIS water-quality products and FUME results were extracted along a transect (yellow line in Fig. 5) perpendicular to the coast. The transect starts at the Match-up point in the Wadden Sea (red circle) and ends in the Central North Sea. The results are shown in the three panels of Fig. 9. Within the first 30 pixels the waters are within or close to the Wadden Sea, characterised by very high loads of sediment ($> 1 \text{ g m}^{-3}$) and Yellow Substance (absorption $> 1 \text{ m}^{-1}$ at 442 nm), corresponding to FU values above 7. In the next part (pixels 30–170), the FU-values show a gradual gradient from 7 to 3, reflecting a decrease in algal pigments (both Algal-1 and Algal-2 products) because both YS and SPM are rather constant in this interval. An interesting feature can be observed in the pixels (170–300) where the Case-2 water algorithm seems to fail (unrealistic high SPM and YS), likely due to additional scatter by cirrus clouds. Fortunately, the FU-scale seems robust and corresponds rather well with the algal-1 product.

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Based on field measurements, a comparison between observed and modelled FU numbers was made (Fig. 10). The correlation between observed and modelled FU numbers is around the 1 : 1 line. However, full spectral derived FU correlates better ($R^2 = 0.81$) with in-situ data than MERIS derived FU ($R^2 = 0.74$). More spectral information provides a more reliable FU estimate. The largest outliers were found in the 11–17 FU range. These are the green-yellowish water colours, which corresponds to ~ 500–600 nm of the visible light. This is also the part of the light spectrum with is covered by only one MERIS band (560 nm) which leads to the largest loss of information.

Comparing the MERIS winter FU maps of the Northern and Southern Red Sea of Fig. 6 with the winter IDW FU map of Fig. 11 similar patterns can be recognized despite a time cap of over a century between data acquisition. When we compare these maps it seems that the colour of the Red Sea did not change significantly over time, although we cannot say anything about intermediate colour changes between 1899 and 2002. Within the red circles the MERIS FU map (Fig. 6) shows FU2 for the northern location and FU8 for the southern location, while the IDW FU map gives identical values at both locations.

The FU map for the Yellow Sea, based on 2882 FU in-situ observations collected during the winters of 1930 to 1999, is shown in Fig. 12. The Yellow Sea shows FU numbers of FU4 at open sea areas to values of FU20 in front of the outflow of the Yangtze River. Both, the MERIS map of Fig. 7 and the IDW interpolation of Fig. 12 show similar colour patterns. The red circles in the MERIS map indicate FU9 in the lower and FU11 in the upper red circle. In the IDW map respectively FU12 and FU14 are indicated. A possible explanation of the bluing of the water (bluer colours show up in the 2008/9 map) is the reduced outflow of Yangtze water into the Yellow Sea due to the hydroelectric Three Gorges Dam which became operational in 2003. The effect is a reduced upwelling and thus productivity resulting in less green water (Chen, 2000; Gong et al., 2006).

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Discussion and conclusions

In this paper an algorithm is presented that allows retrieval of the Forel–Ule sea colour from the MERIS satellite sensor. The Forel–Ule colour can be seen as the colour standard closest to the real colour of water. The elegance of our algorithm is that it converts multispectral observations to one simple number that is only dependent on a well known universal set of colourimetric functions. The classification of sea water is simplified by means of a numerical value between 1 and 21, instead of a classification by a normalized water-leaving spectral reflectance signature, or the concentrations of the dominant optical constituents.

The approach is demonstrated by the processing of multispectral observations of oceans and coastal waters made by the MERIS ocean colour sensor, to FU maps that cover colour classes between indigo blue, green and brown. Five different seas were selected worldwide; these were processed to obtain FU maps. The maps show very detailed patterns and gradients, mainly in the near coastal zone as expected by the more outspoken hydrographical gradients there. When the MERIS maps of sea and ocean colour distribution were compared with ground truth Forel–Ule observations mapped in the same season, similar patterns and FU numbers were observed, even when FU’s of more than a century ago were processed. This opens new ways to study the spatial and temporal evolution of the colour of the sea worldwide. The FUME algorithm can easily be adapted to data from other satellites which have enough bands in the visible part of the spectrum to properly derive the colour of the water.

Acknowledgements. Marieke Eleveld, Steef Peters and Reinold Pasterkamp from the Institute for Environmental Studies, Free University, Amsterdam are thanked for their initiative to develop MATLAB routines that were used and adapted to generate satellite derived Forel–Ule maps. Menno Regeling is thanked for the Forel–Ule data collection in the North Sea (June 2001). Thanks are due to W. Gieskes (Dept. Ocean Ecosystems, University of Groningen, The Netherlands) for discussions and comments. MERIS data has been provided by the European Space Agency (ESA).

MERIS-based Forel–Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Antoine, D., Morel, A., Gordon, H. R., Banzon, V. F., and Evans, R. H.: Bridging ocean colour observations of the 1980s and 2000s in search of long-term trends, *J. Geophys. Res.* 110, 22 pp, C06009, doi:10.1029/2004JC002620, 2005.
- 5 Behrenfeld, M. J., O'Malley, R. T., Siegel, D. A., McClain, C. R., Sarmiento, J. L., Feldman, G. C., Milligan, A. J., Falkowski, P. G., Letelier, R. M., and Boss, E. S.: Climate-driven trends in contemporary ocean productivity, *Nature*, 444, 752–755, 2006.
- Boyer, T. P., Antonov, J. I., Garcia, H. E., Johnson, D. R., Locarnini, R. A., Mishonov, A. V., Pitcher, M. T., Baranova, O. K., and Smolyar, I. V.: World ocean database 2005, in: NOAA
- 10 Atlas NESDIS 60, edited by: Levitus, S., US Government Printing Office, Washington, DC, 190 pp., DVDs, 2006.
- Brockmann, C., Block, T., Faber, O., Fokken, L., and Kock, O.: MERCI – MERIS Catalogue and Inventory, MERIS-ATSR workshop. ESA, ESRIN, 2005.
- Chen, C.-T. A.: The Three Gorges dam: reducing the upwelling and thus productivity in the East
- 15 China Sea, *Geophys. Res. Lett.*, 27, 381–383, 2000.
- CIE: Commission Internationale de l'Éclairage proceedings, 1931, Cambridge University Press, 1932.
- ESA: MERIS Algorithm Theoretical Basis Documents. <http://envisat.esa.int/instruments/meris/pdf/>, last access: 7 January 2012.
- 20 Forel, F. A.: Une nouvelle forme de la gamme de couleur pour l'étude de l'eau des lacs, in: Archives des Sciences Physiques et Naturelles/Société de Physique et d'Histoire Naturelle de Genève, VI, 25 pp., 1890.
- Gong, G.-C., Chang, J., Chiang, K.-P., Hsiung, T.-M., Hung, C.-C., Duan, S.-W., and Codispoti, L. A.: Reduction of primary production and changing of nutrient ratio in the East
- 25 China Sea: effect of the Three Gorges dam?, *Geophys. Res. Lett.*, 33, 4 pp, L14609, doi:10.1029/2006GL025800, 2006.
- Gordon, H. R. and Voss, K. J.: MODIS Normalized Water-leaving Radiance Algorithm Theoretical Basis Document, Vol. MOD 18, Version 4, 1999.
- Heuermann, R., Reuter, R., and Willkomm, R.: RAMSES, a modular multispectral radiometer for light measurements in the UV and VIS, *SPIE Proc. Ser.*, 3821, 279–285, 1999.
- 30 Hommersom, A., Peters, S., Wernand, M. R., and De Boer, J.: Spatial and temporal variability in bio-optical properties of the Wadden Sea, *Estuar. Coast. Shelf S.*, 83, 360–370, 2009.

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- IOCCG: Minimum Requirement for an Operational Ocean-Colour Sensor for the Open Ocean, report number 1, <http://www.ioccg.org/reports.ioccg.html>, last access 7 January 2012, 1998.
- Lee, Z. P., Carder, K. L., Hawes, S. K., Steward, R. G., Peacock, T. G., and Davis, C. O.: Model for the interpretation of hyperspectral remote-sensing reflectance, *Appl. Optics*, 33, 5721–5732, 1994.
- Mobley, C. D.: *Light and Water: Radiative Transfer in Natural Waters*, Academic Press, San Diego, 1994.
- Mueller, J. L., Davis, C., Arnone, R., Frouin, R., Carder, K., Lee, Z. P., Steward, R. G., Hooker, S., Mobley, C. D., and McLean, S.: Above-Water Radiance and Remote Sensing Reflectance Measurement and Analysis Protocols, *Ocean Optics Protocols for Satellite Ocean Colour Sensor Validation*, Revision 4, Volume III: Radiometric Measurements and Data Analysis Protocols, NASA/TM-2003-21621/Rev4 III, 21–31, 2003.
- Odermat, D., Gitelson, A., Brando, V. E., and Schaepman, M.: Review of constituent retrieval in optically deep and complex waters from satellite imagery, *Remote Sens. Environ.*, 118, 116–126, 2012.
- Polovina J. J., Howell, E. A., and Abecassis, M.: Ocean's least productive waters are expanding, *Geophys. Res. Lett.*, 35, 5 pp., L03618, doi:10.1029/2007GL031745, 2008.
- Rast, M., Bezy, J. L., and Bruzzi, S.: The ESA Medium Resolution Imaging Spectrometer MERIS: a review of the instrument and its mission, *Int. J. Remote Sens.*, 20, 1681–1702, 1999.
- Tilstone, G., Peters, S. W. M., Van der Woerd, H., Eleveld, M., Ruddick, K., Schoenfeld, W., Krasemann, H., Martinez-Vicente, V., Blondeau-Patissier, D., Röttgers, R., Sørensen, K., Jorgensen, P., and Shutler, J.: Variability in specific-absorption properties and their use in a semi-analytical ocean colour algorithm for MERIS in North Sea and Western English Channel coastal waters, *Remote Sens. Environ.*, 118, 320–338, 2012.
- Ule, W.: Die bestimmung der Wasserfarbe in den Seen, *Kleinere Mittheilungen*, in: Dr. A. Petermanns Mittheilungen aus Justus Perthes geographischer Anstalt, Justus Perthes, Gotha, 70–71, 1892.
- Van der Woerd, H. J. and Pasterkamp, R.: HYDROPT: a fast and flexible method to retrieve chlorophyll *a* from multispectral satellite observations of optically complex coastal waters, *Remote Sens. Environ.*, 112, 1795–1807, 2008.
- Watson, D. F. and Philip, G. M.: A refinement of inverse distance weighted interpolation, *Geo-processing*, 2, 315–327, 1985.

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Wernand, M. R. and Gieskes, W. W. W.: Ocean optics from 1600 (Hudson) to 1930 (Raman); Shift in interpretation of natural water colouring, Union des océanographes de France, 2011.
- Wernand, M. R. and Van der Woerd, H. J.: Spectral analysis of the Forel–Ule ocean colour comparator scale, J. Eur. Opt. Soc.-Rapid, 5, 7 pp, 10014s, doi:10.2971/jeos.2010.10014s, 2010a.
- Wernand, M. R. and Van der Woerd, H. J.: Ocean colour changes in the North Pacific since 1930, J. Eur. Opt. Soc.-Rapid, 5, 6 pp, 10015s, doi:10.2971/jeos.2010.10015s, 2010b.
- Wernand, M. R., van der Woerd, H. J., and Gieskes, W. W. W.: Trends in ocean colour and chlorophyll concentration from 1889 to 2000, worldwide, Prog. Oceanogr., under review, 2012.
- Wyszecki, G. G. and Stiles, W. S.: Colour Science; Concepts and Methods, Quantitative Data and Formulae, 2nd edition, Wiley classics Library edition, Wiley & sons, inc., 127–129, 2000.

MERIS-based Forel–Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**MERIS-based
Forel-Ule
classification**

M. R. Wernand et al.

Table 1. Central wavelengths of the first nine MERIS spectral bands. All bands have a width of 10 nm, with exception of band 8 (7.5 nm).

MERIS band	Wavelength (nm)	MERIS band	Wavelength (nm)
1	412.5	6	620
2	442.5	7	665
3	490	8	681.25
4	510	9	708
5	560		

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**MERIS-based
Forel-Ule
classification**

M. R. Wernand et al.

Table 2. Chromaticity coordinates (x , y) of the FU scale numbers as determined by Wernand and Van der Woerd (2010a).

No.	x	y	No.	x	y	No.	x	y
FU1	0.189	0.161	FU8	0.311	0.439	FU15	0.410	0.478
FU2	0.196	0.194	FU9	0.337	0.463	FU16	0.418	0.472
FU3	0.213	0.255	FU10	0.363	0.480	FU17	0.427	0.466
FU4	0.229	0.301	FU11	0.388	0.490	FU18	0.440	0.458
FU5	0.242	0.331	FU12	0.394	0.488	FU19	0.453	0.448
FU6	0.263	0.373	FU13	0.397	0.486	FU20	0.462	0.440
FU7	0.290	0.415	FU14	0.404	0.482	FU21	0.473	0.429

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**MERIS-based
Forel-Ule
classification**

M. R. Wernand et al.

Table 3. Angles α_i of FU the number in degrees ($i = 1$ to 21) and the 20 boundary angles α_{iT} that are used in the discrete classification of ocean colour.

i	α_i^0	α_{iT}^0	If $\alpha_M > \alpha_{iT}$ then FU =	i	α_i^0	α_{iT}^0	If $\alpha_M > \alpha_{iT}$ then FU =
1	229.94	227.68	1	12	68.49	67.93	12
2	225.41	219.27	2	13	67.36	65.98	13
3	213.13	205.19	3	14	64.60	63.35	14
4	197.25	189.20	4	15	62.11	60.37	15
5	181.15	165.71	5	16	58.62	56.64	16
6	150.26	133.96	6	17	54.65	52.09	17
7	117.66	109.85	7	18	49.53	46.75	18
8	102.05	95.14	8	19	43.96	41.82	19
9	88.24	83.38	9	20	39.67	36.98	20
10	78.53	74.62	10	21	34.28		21
11	70.71	69.60	11				

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**MERIS-based
Forel-Ule
classification**

M. R. Wernand et al.

Table 4. Reference to the acquired MERIS Reduced Resolution images.

Sea	Area	Product name	Start time UTC
North Sea	1	MER_RR_2PQBCM20060504	4 May 2006 10:11:24
Red Sea north	2	MER_RR_2PQBCM20031222	22 Dec 2003 07:56:09
Red Sea south	3	MER_RR_2PQBCM20031223	23 Dec 2003 07:26:02
Yellow Sea	4	MER_RR_2PPBCM20090211	11 Feb 2009 02:27:39
Sea of Japan	5	MER_RR_2PQBCM20040614	14 Jun 2004 01:07:24

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**MERIS-based
Forel-Ule
classification**

M. R. Wernand et al.

Table 5. RGB values for the reproduction of the FU-legend.

FU nr.	R	G	B	FU nr.	R	G	B
1	33	88	188	12	148	182	96
2	49	109	197	13	165	188	118
3	50	124	187	14	170	184	109
4	75	128	160	15	173	181	95
5	86	143	150	16	168	169	101
6	109	146	152	17	174	159	92
7	105	140	134	18	179	160	83
8	117	158	114	19	175	138	68
9	123	166	84	20	164	105	5
10	125	174	56	21	161	77	4
11	149	182	69				

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MERIS-based Forel-Ule classification

M. R. Wernand et al.




Fig. 1. The CIE 1931 2° Colour Matching Functions for \bar{x} (red), \bar{y} (green) and \bar{z} (blue) determined per nanometre. The black line shows a reconstructed spectrum (Yellow Sea) of MERIS reflectance measured at nine bands (open circles).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 2. The chromaticity coordinates, based upon transmission measurements, of the FU scale colours 1 to 21 (black circles) and the white point W where $x_W = y_W = 1/3$. The outer curved boundary is the spectral locus, with the corresponding mono-chromatic wavelengths shown in nanometre.

MERIS-based Forel-Ule classification

M. R. Wernand et al.




Fig. 3. Chromaticity diagram with scale colours FU1 to FU21 shown in dots, relative to the white point that is set at the origin. As an example the angle α_i (102.05° , see Table 3), determining the position of FU8, is given.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Areas where MERIS data were extracted from ESA's MERCI database.

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 5. The spring FU map of the turbid North Sea (4 May 2006) shows FU values ranging from 3 to 9 (red circle near England FU6). The Wadden Sea area within barrier islands north of Holland shows values between FU9 and FU18. The area within the red circle indicates a pixel value of FU = 14. The yellow line shows the transect extracted from the MERIS image and shown in Fig. 9.




Fig. 6. The winter FU map of the Red Sea, dated 22 (left) and 23 December 2003 (right) shows that open water of the northern part is mainly bluish FU1 to FU2 (in red circle FU2) and near coast values are around FU3. The southern part shows more greenish coloured water (in red circle FU = 8).

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. The winter FU map of the Yellow Sea, acquisition date 11 February 2009 (left) and the summer FU map of the Sea of Japan, acquisition date 14 June 2004 (right). The Yellow Sea is mainly greenish brown with FU7 up to FU17 (FU9 in the lower red circle and FU11 in the upper red circle). The Sea of Japan, a “blue sea”, shows summer values of around FU2 to FU3 (in red circle FU2). East of Hokkaido phytoplankton abundance greens the water to FU10 (FU9 within the red circle east).

MERIS-based Forel-Ule classification

M. R. Wernand et al.




Fig. 8. Example of the pixel values of MERIS normalized water-leaving reflectance spectra of the North Sea (NS), Wadden Sea (WS), Yellow Sea (YS), Sea of Japan (SJP) and the Northern and Southern Red Sea (resp. RS1 and RS2). Notice the similarity in the spectral shapes of the Wadden Sea spectra; WS is a MERIS normalized water-leaving reflectance spectrum and WS (GT) is the ground-truth reflectance spectrum from ship-based measurements.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 9. Transect from the inner Wadden Sea to the Central North Sea. The upper panel shows the FU values, the middle panel the two MERIS products for chlorophyll *a* and the low panel shows the concentration of SPM (units in g m^{-3}) and yellow substance (absorption in m^{-1} at 442 nm)




Fig. 10. Scatterplot of the *FU* numbers modelled using full reflectance spectra (black dots) with trend line (full) and from the reflectance at MERIS bands (circles) with trend line (intermittent) versus in situ observed *FU* numbers.




Fig. 11. The Red Sea map based on 52 FU in-situ observations collected during the winters of 1895 to 1898. Within the northern red circle FU = 2 and within the southern red circle FU = 7. This colour boundary can also be observed 100 yr later in the MERIS map of 6.

MERIS-based Forel-Ule classification

M. R. Wernand et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



